

within the H-burning shell is $B_r \approx 1.5 \times 10^7$ G, although we cannot rule out the presence of stronger fields away from the H-burning shell. This large field strength may indicate that KIC8561221 is the descendant of a magnetic A-type star whose internal field was much stronger than the typical surface fields of $B \sim 3$ kG of A-type stars (19).

In principle, it is possible that another symmetry-breaking mechanism within the core could suppress dipole mode amplitudes. The only other plausible candidate is rapid core rotation. In order for rotation to strongly modify the incoming waves so that they will be trapped in the core, the core must rotate at a frequency comparable to v_{\max} , roughly two orders of magnitude faster than the values commonly measured in red giant cores (2, 6, 20). The depressed dipole mode star KIC8561221 (8) does not exhibit rapid core rotation and disfavors the rotation scenario.

A magnetic field of amplitude $B > 10^4$ G (Fig. 4) could be present in the core of a red giant if it was retained from previous phases of stellar formation and evolution (12). These strong fields may reside within the inner core with little external manifestation apart from the reduced visibility of the dipole modes. However, fields of similar amplitude have been discussed in order to explain the suppression of thermohaline mixing in a small fraction of red giant stars, as inferred from the observations of their surface abundances (21). The inferred core field strength of $B_r \geq 1.5 \times 10^7$ G in KIC8561221 shows that very strong magnetic fields ($B \gg 10^6$ G) can exist within the radiative cores of early RGB stars. Because these fields are probably inherited from previous stages of stellar evolution, slightly weaker ($B \gg 10^5$ G) fields could exist in the cores of exceptional very highly magnetized main sequence stars.

REFERENCES AND NOTES

1. T. R. Bedding, *Solar-like Oscillations: An Observational Perspective* (Cambridge Univ. Press, Cambridge, 2014).
2. P. G. Beck et al., *Science* **332**, 205 (2011).
3. T. R. Bedding et al., *Nature* **471**, 608–611 (2011).
4. B. Mosser et al., *Astron. Astrophys.* **572**, L5 (2014).
5. P. G. Beck et al., *Nature* **481**, 55–57 (2011).
6. B. Mosser et al., *Astron. Astrophys.* **548**, A10 (2012).
7. B. Mosser et al., *Astron. Astrophys.* **537**, A30 (2012).
8. R. A. García et al., *Astron. Astrophys.* **563**, A84 (2014).
9. P. Goldreich, D. A. Keeley, *Astrophys. J.* **212**, 243 (1977).
10. M.-A. Dupret et al., *Astron. Astrophys.* **506**, 57–67 (2009).
11. O. Benomar et al., *Astrophys. J.* **781**, L29 (2014).
12. See the supplementary materials on Science Online.
13. W. J. Chaplin, A. Miglio, *Annu. Rev. Astron. Astrophys.* **51**, 353–392 (2013).
14. E. Corsaro et al., *Astrophys. J.* **757**, 190 (2012).
15. M. Grosjean et al., *Astron. Astrophys.* **572**, A11 (2014).
16. E. Corsaro, J. De Ridder, R. A. García, *Astron. Astrophys.* **579**, A83 (2015).
17. T. R. Bedding et al., *Astrophys. J.* **713**, L176–L181 (2010).
18. T. M. Brown, R. L. Gilliland, R. W. Noyes, L. W. Ramsey, *Astrophys. J.* **368**, 599 (1991).
19. M. Aurière et al., *Astron. Astrophys.* **475**, 1053–1065 (2007).
20. S. Deheuvels et al., *Astron. Astrophys.* **564**, A27 (2014).
21. C. Charbonnel, J.-P. Zahn, *Astron. Astrophys.* **476**, L29–L32 (2007).
22. J. Ballot, C. Barban, C. V. Veer-Menneret, *Astron. Astrophys.* **531**, A124 (2011).

ACKNOWLEDGMENTS

The authors of this paper thank the Kavli Institute for Theoretical Physics and the organizers of the Galactic Archaeology and Precision Stellar Astrophysics program held from January to April 2015. J.F. acknowledges partial support from NSF under grant no. AST-1205732 and through a Lee DuBridge Fellowship at the California Institute of Technology. R.A.G. acknowledge the support of the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement no. 312844 (SPACEINN), and from the Centre National d'Études Spatiales. D.S. acknowledges support from the Australian Research Council. This project was supported by NASA under Theory and

Computational Astrophysics Network grant no. NNX14AB53G and NSF under grants PHY 11-25915 and AST 11-09174.

SUPPLEMENTARY MATERIALS

www.sciencemag.org/content/350/6259/423/suppl/DC1
Supplementary Text
Figs. S1 to S3
Table S1
References (23–56)

29 May 2015; accepted 15 September 2015
10.1126/science.aac6933

PLANT DEVELOPMENT

Transcriptional control of tissue formation throughout root development

Miguel A. Moreno-Risueno,¹ Rosangela Sozzani,^{2*} Galip Gürkan Yardımcı,^{2†} Jalean J. Petricka,^{2‡} Teva Vernoux,³ Ikram Biloul,⁴ Jose Alonso,⁵ Cara M. Winter,² Uwe Ohler,⁶ Ben Scheres,⁴ Philip N. Benfey^{2§}

Tissue patterns are dynamically maintained. Continuous formation of plant tissues during postembryonic growth requires asymmetric divisions and the specification of cell lineages. We show that the BIRDS and SCARECROW regulate lineage identity, positional signals, patterning, and formative divisions throughout *Arabidopsis* root growth. These transcription factors are postembryonic determinants of the ground tissue stem cells and their lineage. Upon further activation by the positional signal SHORT-ROOT (a mobile transcription factor), they direct asymmetric cell divisions and patterning of cell types. The BIRDS and SCARECROW with SHORT-ROOT organize tissue patterns at all formative steps during growth, ensuring developmental plasticity.

Organs are formed, patterned, and maintained during growth. In the root of *Arabidopsis*, tissues are organized as concentric cylinders around the internal vascular tissue. Much progress has been made in identifying factors responsible for patterning some of these tissues, such as the ground tissue. The ground tissue lineage is continuously generated by the cortex endodermis initial stem cell (CEI), which divides in the transverse orientation (anticlinal division) to produce a daughter cell (CEID) and regenerate itself. The ground tissue is patterned when the CEID divides asymmetrically in the longitudinal (periclinal) orientation, gen-

erating two cell types: endodermis and cortex (1). The mobile transcription factor SHORT-ROOT (SHR) moves from the stele to the ground tissue, where SCARECROW (SCR) and the C_2H_2 transcription factor JACKDAW (JKD) sequester it in the nucleus. Nuclear SHR is required for the periclinal asymmetric divisions of the CEID that pattern the ground tissue (2, 3). These divisions are activated through a bistable switch involving SHR, SCR, and other components and correlate with the temporal activation of transcriptional programs (4, 5). Absence of SHR results in abnormal ground tissue patterning, with loss of the endodermis and a remaining single layer of ground tissue due to absence of asymmetric cell divisions (5, 6). Because the ground tissue lineage remains, this indicates that other factors participate in its specification.

Specific roles for JKD and several close relatives have been recently identified (7). JKD and BALD-IBIS regulate SHR movement by promoting its nuclear retention, and cooperatively with MAGPIE (MGP) and NUTCRACKER (NUC) are required for the formative divisions that pattern the ground tissue into cortex and endodermis. Here we show that JKD, MGP, and NUC, along with two new members of this family (collectively known as the BIRDS; table S1) named BLUEJAY (BLJ) and IMPERIAL EAGLE (IME), organize the ground tissue after embryogenesis. They function

¹Department of Biotechnology, Center for Plant Genomics and Biotechnology, Universidad Politécnica de Madrid, 28223 Pozuelo de Alarcón (Madrid), Spain. ²Department of Biology and Howard Hughes Medical Institute, Duke University, Durham, NC 27708, USA. ³Laboratoire de Reproduction et Développement des Plantes, CNRS, INRA, ENS Lyon, UCBL, Université de Lyon, 69364 Lyon, France. ⁴Department of Plant Biology, Wageningen University Research, Wageningen, Netherlands. ⁵Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC 27695, USA. ⁶Berlin Institute for Medical Systems Biology, Max Delbrück Center for Molecular Medicine, 13125 Berlin, Germany. *Present address: Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC 27695, USA. †Present address: Department of Genome Sciences, University of Washington, Seattle, WA 98195, USA. ‡Present address: Carleton College, Northfield, MN 55057, USA. §Corresponding author. E-mail: philip.benfey@duke.edu




Fig. 1. BIRDs are required for ground tissue maintenance and are expressed in the ground tissue. (A to H) Confocal images of roots of wild-type (WT), *shr*, *scr*, *jkd*, *nuc*, *blj*, and *mgp*, at 6 days post-imbibition (dpi), showing the ground tissue (GT) marked by J0571. (I to L) Expression patterns of the BIRDs in WT or *shr* roots at 6 dpi; rec: regulatory regions using recombineering; p: promoter. (M) SCR expression. En, endodermis; C, cortex; QC, quiescent center; LRCEI, lateral root cap/epidermis initial; Pe, pericycle. Scale bars, 10 μ m.




Fig. 2. BLJ, JKD, and SCR specify identity of ground tissue initial cells. (A) Mature embryos of *blj jkd scr*. (B to E) Lineage analyses in *blj jkd scr* using the J0571, cortex (CO2), epidermis (WER), and stele (WOL) markers. Blue arrows: stele; white arrows: epidermis. (F to J) *blj jkd scr* roots from 1 to 5 days after resection (d.a.r.). Regenerated ground tissue (green arrows) is visualized with J0571 marker. Yellow arrows: cells missing J0571 expression. Scale bars, 10 μ m.

as identity determinants of the CEI, which maintains and gives rise to the lineage, and act as effectors of asymmetric cell divisions of the CEID

upon SHR activation. Furthermore, the BIRDs regulate the transcriptional identity of the two ground tissue cell types and form a regulatory

network associated with lineage determination, asymmetric division, cell-type specification, and differentiation.

To explore the role of the BIRDs in ground tissue specification, we used expression from the J0571 enhancer trap line as a robust marker for ground tissue identity (Fig. 1A). This is exemplified by its expression in the mutant layer of *shr* (Fig. 1B). However, its expression was lost in some cells of double- and triple-mutant combinations of BIRDs, and was almost undetectable in the quadruple mutant, *blj jkd mgp nuc* (Fig. 1, D to G), suggesting that these transcription factors play a role in ground tissue identity. We next introgressed combinations of the BIRD mutants into *scr*. In both *jkd scr* and *blj jkd scr*, ground tissue marker expression was reduced, and a number of *blj jkd scr* roots lacked the entire ground tissue (Fig. 1H). These results indicate that the BIRDs and SCR are required for maintenance of ground tissue identity in addition to their established role in patterning divisions and endodermis specification (1, 7).

The mRNAs of *JKD*, *MGP*, and *NUC* are primarily expressed in the ground tissue (6, 7). To accurately determine where the encoded proteins and BLJ and IME accumulate, we tagged the proteins with green fluorescent protein (GFP) (8). BLJ was specifically expressed in the ground tissue, more highly in the shootward part of the meristem, although it was occasionally detected in ground tissue stem cells. Similarly, IME was also expressed more strongly in the shootward part of the meristem. By contrast, *JKD*, *MGP*, and *NUC* expression was higher toward the root tip (fig. S1). Previously, expression of the BIRDs and SCR appeared to require SHR (5, 6). How-

ever, when we introgressed the expression constructs into *shr*, we could detect all of the BIRD fusion proteins, although BLJ expression was reduced in this background (Fig. 1, I to L). SCR is also detectable in the ground tissue of *shr* (9). Taken together, our results indicate that the BIRDs and SCR can be regulated in a manner that is both dependent and independent of SHR, which is consistent with the difference in the phenotypes of *blj jkd scr* and *shr*.

The ground tissue lineage is initially specified in the embryo (10). Inspection of *blj jkd scr* embryos revealed that they developed ground tissue during embryogenesis (Fig. 2A). In seedling roots, we found that the ground tissue was present in mature zones but frequently did not continue to the meristem (fig. S2, A to L). Loss of the ground tissue was detected after germination but became more severe over time (fig. S2M). When the ground tissue disappeared, the epidermis became directly juxtaposed with the stele, as shown by tissue-specific markers (Fig. 2, B to E). These results indicate that the combined activity of the BIRDs and SCR is crucial to maintain the ground tissue lineage postembryonically.

The ground tissue marker J0571 was also lost in cells that were apparently formed by division of ground tissue cells (Fig. 1, D to G) and thus, the critical role of BLJ, JKD, and SCR in maintaining the ground tissue appears to be more than maintenance of the division potential of the CEI. Consistent with this hypothesis, other mutations affecting stem cell niche activity or the orientation plane of niche asymmetric cell divisions do

not result in loss of cell lineages (11, 12). To determine if the loss of ground tissue was due to incorrect specification of the CEI, we regenerated roots from wild-type, *scr*, and *blj jkd scr* mutants after resection of the root tip (fig. S3, A to C). In *blj jkd scr*, there was a low regeneration frequency (fig. S3D), but in those meristems that did regenerate, we found severely impaired regeneration of ground tissue with, at most, one or two cells after 2 days (Fig. 2, F and G). These cells failed to establish a new ground tissue lineage and, although a small amount of division occurred, expression of the ground tissue marker was normally lost (Fig. 2, H to J). *scr* mutants also showed impaired regeneration of the ground tissue lineage and failed to complete an entire layer (fig. S4). Our results indicate that BLJ, JKD, and SCR maintain CEI stem cells and their progeny post-embryonically through specification of CEI identity.

The BIRDs and SCR are involved in regulation of transcription associated with formative divisions of the CEID and ground tissue cells (2, 6, 7). To further investigate the dual role of the BIRDs and SCR in generating and patterning the ground tissue, we reconstructed a gene regulatory network. Among the BIRDs, we focused on BLJ and JKD because they have a specific role in ground tissue establishment and identified their direct targets through chromatin immunoprecipitation (ChIP) followed by sequencing (table S2). To reconstruct the network, we used transcriptionally regulated targets that were identified by the intersection of ChIP-bound genes with previous genome-wide expression data (for SHR), as well




Fig. 3. BIRDs are determinants of ground tissue gene expression and form a network with SHR and SCR. (A) Network of SHR, BIRDs, SCR, and downstream transcription factor hubs. (B) Comparison of transcriptionally regulated targets. (C) Principal component analysis of the transcriptional profiles of ground tissue-expressed genes in different mutants; inset: component weights. (D) Same profiles hierarchically clustered. (E) Redundancy among regulated genes.

as with new microarray data that we generated from *blj jkd scr* root meristems and from ground tissue sorted cells of *shr* mutants in which we amplified BIRDs and SCR expression using the J0571xUAS system [tables S3 and S4; a detailed description of network reconstruction is in (13)]. The resulting network (table S5) showed pronounced overlaps between the genes regulated by more than one transcription factor (Fig. 3A and fig. S5). A more detailed analysis of the network showed that in addition to directly regulating genes (~55%) in the SHR pathway (table S6), the BIRDs and SCR also directly regulate genes in other pathways, suggesting that genes not in the SHR pathway are likely to be involved in maintaining ground tissue identity and may be activated through a transcriptional cascade (Fig. 3B).

To better understand how the BIRDs and SCR regulate ground tissue identity, we analyzed their ability to “rescue” ground tissue gene expression in the *shr* background. As these factors are down-regulated in *shr*, we amplified their expression using the J0571xUAS-driven lines in *shr* (see Methods). We then compared mRNA expression profiles of sorted ground tissue cells from these lines with the expression profile of wild-type ground tissue using principal component analysis (PCA) and hierarchical clustering. As an initial test, we asked if PCA could distinguish between the ground tissue and other root tissues as profiled in the RootMap (14) and found clear

separation (fig. S6, A and B). This provided confidence to use PCA to infer the identity of the transcriptome when each of the BIRDs was expressed in the *shr* ground tissue. Consistently, PCA localized the BIRDs’ ground tissue transcriptomes between *shr* and wild type (Fig. 3C). This was also true for SCR, indicating that a degree of rescue had occurred. In agreement with the PCA, hierarchical clustering separated the ground tissue transcriptome from other root tissues and showed different levels of rescue provided by the BIRDs and SCR (Fig. 3D and fig. S6, C and D). JKD and SCR were able to rescue the ground tissue transcriptome more effectively than did BLJ. This suggests that BLJ’s role may be to regulate a set of genes that are complementary to those regulated by JKD and SCR, leading to a combinatorial action responsible for the phenotype of *blj jkd scr*. The contrasting expression patterns of BLJ and JKD are consistent with complementary activities. In addition, there was greater overlap among genes regulated by SCR and the BIRDs that showed a higher level of rescue of the ground tissue transcriptome than with the BIRDs that showed a lower level of rescue (Fig. 3E).

Our network indicated that SCR, BLJ, JKD, MGP, and NUC are direct targets of SHR, whereas IME is regulated through an intermediate transcription factor. SCR, NUC, and MGP are activated upon SHR induction (5). To better understand the role of BLJ, JKD, and IME downstream of SHR, we

performed a real-time polymerase chain reaction analysis of gene expression after SHR induction (Fig. 4A). We found that mRNA levels of SCR, BLJ, JKD, MGP, NUC, and IME genes peaked at 6 hours, coincident with the time of onset of asymmetric divisions patterning the ground tissue. A large proportion of the genes down-regulated in *shr* are activated by the BIRDs and SCR (Fig. 4B), and some of these had been previously identified to be activated by the BIRDs and SCR (Fig. 4B), and some of these had been previously identified to be activated at the time of the asymmetric divisions (fig. S7A). This indicates that SHR activates the BIRDs and SCR to induce gene expression associated with patterning. It is, therefore, possible that activation of downstream targets could be dependent on the level of SCR and the BIRDs, with SHR being the amplifying signal required for patterning. Inspection of the ground tissue in the J0571xUAS lines showed formative divisions (fig. S7, B to H) with high penetrance when BLJ was expressed in the *shr* background (80% of roots, $n = 20$), whereas other BIRDs and SCR could induce these divisions only with substantially lower penetrance (IME: 50%; others: 10%; $n = 20$).

As suggested by our network, we next asked if BLJ, JKD, and SCR are required for SHR function in the ground tissue. For this purpose, we fused SHR to a nuclear localization signal and expressed it directly in the ground tissue under a two-component system driven by the En7 promoter, which is specific to the endodermis and CEI (fig. S7, I to L). When expressed in a wild-type




Fig. 4. BIRDs are activated by SHR and specify endodermis and cortex. (A) BIRD expression in ground tissue cells at different times after SHR induction. (B) Venn diagrams comparing genes activated by BIRDs and SHR. (C) Heatmap of BIRD contribution to tissue-enriched expression. Ph: phloem; Xy: xylem; Va: vasculature; Col: columella. (D to F) Optical root sections at 10 dpi. White arrows: Casparian strip; green arrows: lignin in xylem. (G to H) Cortex marker in 6 dpi roots. (I and J) Model of postembryonic formation and maintenance of ground tissue. Scale bars, 10 μ m.

background, SHR generated additional ground tissue layers, as previously described (7, 9). However, when expressed in combinations of mutants involving *blj jkd* and *scr*, SHR failed to rescue the formative cell divisions within the ground tissue. These results indicate that BLJ, JKD, and SCR are essential for SHR to carry out ground tissue patterning. Furthermore, analysis of the contribution of the BIRDs and SCR to generate specific gene expression patterns showed that these transcription factors were able to activate expression of endodermis and cortex genes (Fig. 4C). Staining for endodermis-specific attributes (the Casparian strip) in the *shr* J0571xUAS lines showed that BLJ (Fig. 4, D to F), along with the other BIRDs and SCR (fig. S7, M to Q), could induce Casparian strip formation subsequent to periclinal divisions of the ground tissue. Expression of cortex-specific markers required at least JKD, MGP, and NUC (Fig. 4, G and H). SCZ, which is required for expression of some cortex-specific markers (15), is also a target in the network. Our analysis suggests that cortex identity requires multiple inputs from the BIRDs. Therefore, the BIRDs and SCR, in addition to mediating SHR transcriptional competence (7), are endogenous effectors of ground tissue patterning and can provide all the necessary information for the asymmetric divisions that are activated by SHR to pattern the ground tissue.

Cell fate choices in all multicellular organisms are governed by transcription factors. Their combinatorial expression and interactions are key to tissue identity. The BIRDs and SCR play critical roles in maintaining ground tissue identity in postembryonic roots by specifying the CEI stem cells that generate the ground tissue lineage (Fig. 4I). In addition, they are effectors of asymmetric divisions that pattern the progeny of the CEIs (Fig. 4J). The continuous control of multiple steps of tissue formation by the same set of transcription factors, independently of and dependent on positional cues, is a sophisticated mechanism ensuring plasticity in the regulation of cell fate.

REFERENCES AND NOTES

- J. J. Petricka, J. M. Van Norman, P. N. Benfey, *Cold Spring Harb. Perspect. Biol.* **1**, a000497 (2009).
- D. Welch *et al.*, *Genes Dev.* **21**, 2196–2204 (2007).
- H. Cui *et al.*, *Science* **316**, 421–425 (2007).
- A. Cruz-Ramirez *et al.*, *Cell* **150**, 1002–1015 (2012).
- R. Sozzani *et al.*, *Nature* **466**, 128–132 (2010).
- M. P. Levesque *et al.*, *PLoS Biol.* **4**, e143 (2006).
- Y. Long *et al.*, *Plant Cell* **27**, 1185–1199 (2015).
- R. Zhou, L. M. Benavente, A. N. Stepanova, J. M. Alonso, *Plant J.* **66**, 712–723 (2011).
- Y. Helariutta *et al.*, *Cell* **101**, 555–567 (2000).
- M. Pernas, E. Ryan, L. Dolan, *Curr. Biol.* **20**, 818–823 (2010).
- M. Aida *et al.*, *Cell* **119**, 109–120 (2004).
- V. Willemsen *et al.*, *Dev. Cell* **15**, 913–922 (2008).
- Materials and methods are available as supplementary materials on Science Online.
- S. M. Brady *et al.*, *Science* **318**, 801–806 (2007).
- C. A. ten Hove *et al.*, *Curr. Biol.* **20**, 452–457 (2010).

ACKNOWLEDGMENTS

This work was supported by grants from the NIH (R01-GM043778) and the Gordon and Betty Moore Foundation (GBMF3405) to P.N.B.; from the NSF (IOS-1021619-002) to U.O. and P.N.B.; and from MINECO, FEDER/EFDR (BFU2013-41160-P), and FP7 (PCIG11-GA-2012-322082) to M.A.M.-R. M.A.M.-R. is supported by the Ramon y Cajal program (MINECO). C.M.W. by a NRSA F32 GM106690-01 fellowship, I.B. by a NWO VIDI grant, and

B.S. by an ERC Advanced Grant SysArc. J.J.P. was supported by a NIH Ruth L. Kirschstein NRSA F32 GM086976 fellowship I.B. by a NWO VIDI grant, and B.S. by an ERC Advanced Grant SysArc. We thank D. R. McClay, D. M. Pernas, and Y. Long for critical reading of the manuscript. Additional data are in the supplementary materials. The National Center for Biotechnology Information Gene Expression Omnibus accession numbers are GSE60157 (microarrays) and GSE60011 (ChIP-Seq). The authors declare that they have no competing interests.

SUPPLEMENTARY MATERIALS

www.sciencemag.org/content/350/6259/426/suppl/DC1
Materials and Methods
Figs. S1 to S7
Table S1 to S6
References (16–30)

30 July 2015; accepted 22 September 2015
10.1126/science.121171

ALZHEIMER'S DISEASE

Reduced grid-cell-like representations in adults at genetic risk for Alzheimer's disease

Lukas Kunz,^{1,2} Tobias Navarro Schröder,³ Hweeling Lee,¹ Christian Montag,⁴ Bernd Lachmann,⁴ Rayna Sariyska,⁴ Martin Reuter,^{5,6} Rüdiger Stirnberg,¹ Tony Stöcker,¹ Paul Christian Messing-Floeter,^{1,2} Juergen Fell,² Christian F. Doeller,^{3*} Nikolai Axmacher^{1,2,7*†}

Alzheimer's disease (AD) manifests with memory loss and spatial disorientation. AD pathology starts in the entorhinal cortex, making it likely that local neural correlates of spatial navigation, particularly grid cells, are impaired. Grid-cell-like representations in humans can be measured using functional magnetic resonance imaging. We found that young adults at genetic risk for AD (*APOE-ε4* carriers) exhibit reduced grid-cell-like representations and altered navigational behavior in a virtual arena. Both changes were associated with impaired spatial memory performance. Reduced grid-cell-like representations were also related to increased hippocampal activity, potentially reflecting compensatory mechanisms that prevent overt spatial memory impairment in *APOE-ε4* carriers. Our results provide evidence of behaviorally relevant entorhinal dysfunction in humans at genetic risk for AD, decades before potential disease onset.

Late-onset AD is the most common form of dementia and one of the most challenging diseases of modern society (1). Curative therapies are still lacking, presumably because they start too late (2). Therefore, the elucidation of early pathomechanisms underlying symptoms of AD is of high interest. We aimed at identifying one of the potentially earliest neurocognitive pathomechanisms in the development of AD symptoms: We hypothesized entorhinal dysfunction in young *APOE-ε4* carriers. Our hypothesis was built on three previous findings: First, the $\epsilon 4$ allele of the *APOE* gene is the strongest genetic risk factor for late-onset AD (3). Individuals carrying one *APOE-ε4* allele are at threefold increased risk of AD, and those carrying two *APOE-ε4* alleles are at more than 10-fold increased risk (4). Second, early AD histopathology appears in the entorhinal cortex (EC) (5), where tau abnormal-

ities can already be observed in adults under the age of 30 (6), especially in *APOE-ε4* carriers (7). Third, the EC contains grid cells, a cell type involved in spatial navigation. Grid cells fire whenever animals (8) or humans (9) traverse the vertices of an internally generated grid tiling the spatial environment into equilateral triangles. Their function has been linked to path integration (10, 11), error correction (12), and the maintenance of place cells (13), which exhibit only a singular firing field (14). Hence, a possible dysfunction of grid cells may provide an explanation for the symptom of spatial disorientation in patients suffering from AD. Proxies for grid cells, termed grid-cell-like representations, are detectable in humans by functional magnetic resonance imaging (fMRI). The blood oxygenation level-dependent signal of the EC depends on movement direction with sixfold rotational symmetry. More specifically, the contrast of brain activity during movements aligned versus misaligned to the main axes of a putative grid in a virtual arena leads to a macroscopically visible fMRI signal in the right EC (15).

We examined the effect of *APOE-ε4* on grid-cell-like representations by comparing two groups of healthy young adults ($n = 38$ *APOE-ε4/ε3* carriers, termed "risk participants" from now on; $n = 37$ $\epsilon 3/\epsilon 3$ carriers, "control participants"; table S1). Participants completed a previously established

¹German Center for Neurodegenerative Diseases (DZNE), Bonn, Germany. ²Department of Epileptology, University of Bonn, Bonn, Germany. ³Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, Netherlands. ⁴Department of Psychology, Ulm University, Ulm, Germany. ⁵Department of Psychology, University of Bonn, Bonn, Germany. ⁶Center for Economics and Neuroscience, University of Bonn, Bonn, Germany. ⁷Department of Neuropsychology, Institute of Cognitive Neuroscience, Ruhr-University Bochum, Bochum, Germany. *These authors contributed equally to this work. †Corresponding author. E-mail: nikolai.axmacher@rub.de



Transcriptional control of tissue formation throughout root development

Miguel A. Moreno-Risueno *et al.*

Science **350**, 426 (2015);

DOI: 10.1126/science.aad1171

This copy is for your personal, non-commercial use only.

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

Permission to republish or repurpose articles or portions of articles can be obtained by following the guidelines [here](#).

The following resources related to this article are available online at www.sciencemag.org (this information is current as of October 22, 2015):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/content/350/6259/426.full.html>

Supporting Online Material can be found at:

<http://www.sciencemag.org/content/suppl/2015/10/21/350.6259.426.DC1.html>

This article **cites 29 articles**, 11 of which can be accessed free:

<http://www.sciencemag.org/content/350/6259/426.full.html#ref-list-1>

Downloaded from www.sciencemag.org on October 22, 2015